



- 1 48. (New) A method in accordance with claim 47 wherein B is a phenyl group
2 having one to three substituents selected from the group consisting of -CO₂Me, trifluoromethyl,
3 fluoro, chloro, and methoxy.

- 1 49. (New) A method in accordance with claim 44 having the formula:



- 2 wherein B is a phenyl group optionally substituted with one to three substituents
3 selected from the group consisting of halogen, (C₁-C₄)haloalkyl, (C₁-C₄)haloalkoxy,
4 (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)heteroalkyl, phenyl, phenoxy and -CO₂Me;
5 R¹⁶ and R¹⁷ are independently selected from the group consisting of hydrogen,
6 (C₁-C₈)alkyl and (C₁-C₈)heteroalkyl; or R¹⁶ and R¹⁷ together with the nitrogen atom to which
7 they are attached form a 4- to 7-membered heterocyclic ring optionally having additional
8 heteroatoms as ring members and optionally substituted with substituents selected from the
9 group consisting of (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, hydroxyl, amino, acetoamido and phenyl.
10
- 1 50. (New) A method in accordance with claim 49 wherein -NR¹⁶R¹⁷ is
2 selected from the group consisting of:



55. (New) A composition of claim 52 wherein B is a phenyl group optionally having one to three substituents selected from the group consisting of halogen, (C₁-C₄)haloalkyl, (C₁-C₄)alkyl, (C₁-C₄)alkoxy and -CO₂Me.

56. (New) A composition of claim 55 wherein B is a phenyl group having one to three substituents selected from the group consisting of -CO₂Me, trifluoromethyl, fluoro, chloro, and methoxy.

57. (New) A composition of claim 52 having the formula:

